

COMMISSIONS 27 AND 42 OF THE IAU
INFORMATION BULLETIN ON VARIABLE STARS
Number 4778

Konkoly Observatory
Budapest
13 October 1999
HU ISSN 0374 – 0676

NEW ELEMENTS AND LIGHT CURVE OF CR TAURI

(BAV MITTEILUNGEN NO. 123)

FRANZ AGERER

Bundesdeutsche Arbeitsgemeinschaft für Veränderliche Sterne e.V. (BAV), Munsterdamm 90, D-12169 Berlin, Germany

Name of the object:
CR Tau = S3949 Tau

Observatory and telescope:
Private Observatory, 20-cm SCT

Detector:	SBIG ST6 camera
------------------	-----------------

Filter(s):	None
-------------------	------

Comparison star(s):	GSC 1862.1685
----------------------------	---------------

Check star(s):	GSC 1862.1725
-----------------------	---------------

Transformed to a standard system:	No
--	----

Remarks:
CR Tau was discovered by Hoffmeister (1949a). 15 photographic minima were used to derive the first elements (Hoffmeister 1949b). These elements are listed in the GCVS. Recently we could observe photoelectrically ten new minima, which showed the GCVS-ephemeris to be a spurious one. The minimum times are calculated according to the Kwee-van Woerden method (1956). Using all available photoelectric minima, a weighted least squares fit led to the new ephemeris: $\text{Min I} = \text{HJD } 2451195.4818 \pm 2 + 0^{\text{d}}68270353 \pm 15 \times E.$

Acknowledgements:
Data from the Lichtenknecker Database were used

Table 1: Observed times of minima for CR Tau, epochs and residuals computed with respect to the linear ephemeris derived in this paper.

JD hel. 2400000+	Type*	Epoch	$O - C$	Ref.	JD hel. 2400000+	Type*	Epoch	$O - C$	Ref.
26004.38	P	-36899.0	-0.02	[1]	31449.65	P	-28923.0	+0.00	[1]
26313.70	P	-36446.0	+0.03	[1]	49688.756	E:	-2207.0	+0.001	[2]
26355.27	P	-36385.0	-0.04	[1]	49726.307	E:	-2152.0	+0.003	[2]
26634.51	P	-35976.0	-0.03	[1]	49734.4942	E	-2140.0	-0.0020	[2]
26662.48	P	-35935.0	-0.05	[1]	49756.3431	E	-2108.0	+0.0003	[2]
26718.45	P	-35853.0	-0.06	[1]	50428.4626	E	-1124.5	-0.0018	[2]
26987.51	P	-35459.0	+0.01	[1]	50464.3073	E	-1071.0	+0.0010	[2]
27342.51	P	-34939.0	+0.01	[1]	50849.3518	E	-507.0	+0.0007	[2]
27368.45	P	-34901.0	+0.00	[1]	50863.3450	E	-487.5	-0.0015	[2]
27394.39	P	-34863.0	+0.00	[1]	50864.3710	E	-485.0	+0.0004	[3]
27396.41	P	-34860.0	-0.03	[1]	50864.3711	E	-485.0	+0.0005	[2]
27535.36	P	-34657.5	-0.01	[1]	51185.2412	E	-15.0	-0.0000	[4]
27688.60	P	-34432.0	-0.03	[1]	51195.4821	E	0.0	+0.0003	[2]
31447.63	P	-28926.0	+0.03	[1]					

* P denotes photographic minima (weight 1) and E CCD observed minima (weight 10).

Those marked with ‘:’ got reduced weight (5).

[1]: Hoffmeister (1949b), [2]: Agerer: this paper, [3]: Diethelm (1998), [4]: Diethelm (1999)




Figure 1. Differential light curve of CR Tau, drawn with the new ephemeris

References:

Diethelm, R., 1998, *BBSAG Bulletin* **117**

Diethelm, R., 1999, *BBSAG Bulletin* **119**

Hoffmeister, C., 1949a, *Erg. AN* **12**, 1

Hoffmeister, C., 1949b, *VSS* **1**, 3

Kholopov, P. N. et al. 1985, *General Catalogue of Variable Stars*, 4th Edition, Nauka, Moscow

Kwee, K. K., van Woerden, H., 1956, *Bull. Astr. Inst. Netherlands* **12**, 327